

# Chapter 03.01

## Solution of Quadratic Equations

After reading this chapter, you should be able to:

1. find the solutions of quadratic equations,
2. derive the formula for the solution of quadratic equations,
3. solve simple physical problems involving quadratic equations.

### What are quadratic equations and how do we solve them?

A quadratic equation has the form

$$ax^2 + bx + c = 0, \text{ where } a \neq 0$$

The solution to the above quadratic equation is given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

So the equation has two roots, and depending on the value of the discriminant,  $b^2 - 4ac$ , the equation may have real, complex or repeated roots.

If  $b^2 - 4ac < 0$ , the roots are complex.

If  $b^2 - 4ac > 0$ , the roots are real.

If  $b^2 - 4ac = 0$ , the roots are real and repeated.

### Example 1

Derive the solution to  $ax^2 + bx + c = 0$ .

#### Solution

$$ax^2 + bx + c = 0$$

Dividing both sides by  $a$ , ( $a \neq 0$ ), we get

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0$$

Note if  $a = 0$ , the solution to

$$ax^2 + bx + c = 0$$

is

$$x = -\frac{c}{b}$$

Rewrite

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0$$

as

$$\left(x + \frac{b}{2a}\right)^2 - \frac{b^2}{4a^2} + \frac{c}{a} = 0$$

$$\begin{aligned} \left(x + \frac{b}{2a}\right)^2 &= \frac{b^2}{4a^2} - \frac{c}{a} \\ &= \frac{b^2 - 4ac}{4a^2} \end{aligned}$$

$$\begin{aligned} x + \frac{b}{2a} &= \pm \sqrt{\frac{b^2 - 4ac}{4a^2}} \\ &= \pm \frac{\sqrt{b^2 - 4ac}}{2a} \end{aligned}$$

$$\begin{aligned} x &= -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \end{aligned}$$

### Example 2

A ball is thrown down at 50 mph from the top of a building. The building is 420 feet tall. Derive the equation that would let you find the time the ball takes to reach the ground.

#### Solution

The distance  $s$  covered by the ball is given by

$$s = ut + \frac{1}{2}gt^2$$

where

$u$  = initial velocity (ft/s)

$g$  = acceleration due to gravity (ft/s<sup>2</sup>)

$t$  = time (s)

Given

$$u = 50 \frac{\text{miles}}{\text{hour}} \times \frac{1 \text{ hour}}{3600 \text{ s}} \times \frac{5280 \text{ ft}}{1 \text{ mile}}$$

$$= 73.33 \frac{\text{ft}}{\text{s}}$$

$$g = 32.2 \frac{\text{ft}}{\text{s}^2}$$

$$s = 420 \text{ ft}$$

we have

$$420 = 73.33t + \frac{1}{2}(32.2)t^2$$

$$16.1t^2 + 73.33t - 420 = 0$$

The above equation is a quadratic equation, the solution of which would give the time it would take the ball to reach the ground. The solution of the quadratic equation is

$$t = \frac{-73.33 \pm \sqrt{73.33^2 - 4 \times 16.1 \times (-420)}}{2(16.1)}$$
$$= 3.315, -7.870$$

Since  $t > 0$ , the valid value of time  $t$  is 3.315 s.

---

### NONLINEAR EQUATIONS

---

Topic	Solution of quadratic equations
Summary	Textbook notes on solving quadratic equations
Major	General Engineering
Authors	Autar Kaw
Date	December 23, 2009
Web Site	<a href="http://numericalmethods.eng.usf.edu">http://numericalmethods.eng.usf.edu</a>

---